

ADRA1B Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP9404c**Specification**

ADRA1B Antibody (Center) Blocking Peptide - Product InformationPrimary Accession [P35368](#)**ADRA1B Antibody (Center) Blocking Peptide - Additional Information**

Gene ID 147

Other Names

Alpha-1B adrenergic receptor, Alpha-1B adrenoreceptor, Alpha-1B adrenoceptor, ADRA1B

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

ADRA1B Antibody (Center) Blocking Peptide - Protein Information

Name ADRA1B

Function

This alpha-adrenergic receptor mediates its action by association with G proteins that activate a phosphatidylinositol- calcium second messenger system. Its effect is mediated by G(q) and G(11) proteins. Nuclear ADRA1A-ADRA1B heterooligomers regulate phenylephrine (PE)-stimulated ERK signaling in cardiac myocytes.

Cellular Location

Nucleus membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Cytoplasm Membrane, caveola. Note=Location at the nuclear membrane facilitates heterooligomerization and regulates ERK- mediated signaling in cardiac myocytes. signaling in cardiac myocytes Colocalizes with GNAQ, PLCB1 as well as LAP2 at the nuclear membrane of cardiac myocytes

ADRA1B Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

ADRA1B Antibody (Center) Blocking Peptide - Images

ADRA1B Antibody (Center) Blocking Peptide - Background

Alpha-1-adrenergic receptors (alpha-1-ARs) are members of the G protein-coupled receptor superfamily. They activate mitogenic responses and regulate growth and proliferation of many cells. There are 3 alpha-1-AR subtypes: alpha-1A, -1B and -1D, all of which signal through the Gq/11 family of G-proteins and different subtypes show different patterns of activation. This protein encodes alpha-1B-adrenergic receptor, which induces neoplastic transformation when transfected into NIH 3T3 fibroblasts and other cell lines. Thus, this normal cellular gene is identified as a protooncogene. This protein comprises 2 exons and a single large intron of at least 20 kb that interrupts the coding region.

ADRA1B Antibody (Center) Blocking Peptide - References

Mathias,R.A., J. Allergy Clin. Immunol. 125 (2), 336-346 (2010)Jensen,B.C., Circ Heart Fail 2 (6), 654-663 (2009)Gratacos,M., Am. J. Med. Genet. B Neuropsychiatr. Genet. 150B (6), 808-816 (2009)