

**IFN- $\gamma$**   
Catalog # PVGS1209

## Specification

---

### IFN- $\gamma$ - Product Information

Primary Accession [P01579](#)  
**Species**  
Human

**Sequence**  
Gln24-Gln166

**Purity**  
> 95% as analyzed by SDS-PAGE  
> 95% as analyzed by HPLC

**Endotoxin Level**  
< 0.2 EU/  $\mu$ g of protein by gel clotting method

**Biological Activity**  
ED<sub>50</sub> < 2.0 ng/ml, measured in a cytotoxicity assay using HT-29 cells, corresponding to a specific activity of  $> 5.0 \times 10^5$  units/mg.

**Expression System**  
CHO

Formulation **Lyophilized after extensive dialysis against PBS.**

**Reconstitution**  
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH<sub>2</sub>O or PBS up to 100  $\mu$ g/ml.

**Storage & Stability**  
Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

### IFN- $\gamma$ - Additional Information

**Gene ID** 3458

**Other Names**  
Interferon gamma, IFN-gamma, Immune interferon, IFNG

**Target Background**  
Human Interferon gamma (hIFN- $\gamma$ ) is a macrophage-activating factor and the lone member of Interferon type II. The active form of IFN- $\gamma$  is an antiparallel dimer that interacts with the receptor IFN- $\gamma$ R1 and sets off IFN- $\gamma$ /JAK/STAT pathway. IFN- $\gamma$  signaling does diverse biological functions primarily related to host defense and immune regulation, including antiviral and antibacterial

defense, apoptosis, inflammation, and innate and acquired immunity. While IFN- $\gamma$ -induced inflammatory cascade summons a variety of immune-related cell types, such as macrophages, natural killer (NK) cells and cytotoxic T lymphocytes (CTLs), IFN- $\gamma$  is also implicated in resistance to NK cell and CTL responses and in immune escape in a variety of cancers.

## IFN- $\gamma$ - Protein Information

### Name IFNG

### Function

Type II interferon produced by immune cells such as T-cells and NK cells that plays crucial roles in antimicrobial, antiviral, and antitumor responses by activating effector immune cells and enhancing antigen presentation (PubMed: [16914093](http://www.uniprot.org/citations/16914093)), PubMed: [8666937](http://www.uniprot.org/citations/8666937)). Primarily signals through the JAK-STAT pathway after interaction with its receptor IFNGR1 to affect gene regulation (PubMed: [8349687](http://www.uniprot.org/citations/8349687)). Upon IFNG binding, IFNGR1 intracellular domain opens out to allow association of downstream signaling components JAK2, JAK1 and STAT1, leading to STAT1 activation, nuclear translocation and transcription of IFNG-regulated genes. Many of the induced genes are transcription factors such as IRF1 that are able to further drive regulation of a next wave of transcription (PubMed: [16914093](http://www.uniprot.org/citations/16914093)). Plays a role in class I antigen presentation pathway by inducing a replacement of catalytic proteasome subunits with immunoproteasome subunits (PubMed: [8666937](http://www.uniprot.org/citations/8666937)). In turn, increases the quantity, quality, and repertoire of peptides for class I MHC loading (PubMed: [8163024](http://www.uniprot.org/citations/8163024)). Increases the efficiency of peptide generation also by inducing the expression of activator PA28 that associates with the proteasome and alters its proteolytic cleavage preference (PubMed: [11112687](http://www.uniprot.org/citations/11112687)). Up-regulates as well MHC II complexes on the cell surface by promoting expression of several key molecules such as cathepsins B/CTSB, H/CTSH, and L/CTSL (PubMed: [7729559](http://www.uniprot.org/citations/7729559)). Participates in the regulation of hematopoietic stem cells during development and under homeostatic conditions by affecting their development, quiescence, and differentiation (By similarity).

### Cellular Location

Secreted.

### Tissue Location

Released primarily from activated T lymphocytes.

## IFN- $\gamma$ - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)
- IFN- $\gamma$  - Images**