

**NAP-2/CXCL7**  
**Catalog # PVGS1217****Specification**

---

**NAP-2/CXCL7 - Product Information**

Primary Accession [Q99ME0](#)  
**Species**  
Rat

**Sequence**  
Ile46-Ile107

**Purity**  
> 97% as analyzed by SDS-PAGE<br>> 97% as analyzed by HPLC

**Endotoxin Level**  
< 1 EU/ µg of protein by LAL method

**Biological Activity**  
Fully biologically active when compared to standard. The biologically active determined by a chemotaxis bioassay using human CXCR2 transfected murine BaF3 cells is in a concentration range of 0.1-1.0 ng/ml.

**Expression System**  
E. coli

**Theoretical Molecular Weight**  
6.8 kDa

Formulation **Lyophilized from a 0.2 µm filtered solution in PBS, pH 7.4.**

**Reconstitution**  
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml.

**Storage & Stability**  
Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

**NAP-2/CXCL7 - Additional Information**

**Target Background**  
Neutrophil Activating Peptide 2 (NAP-2) is proteolytically processed carboxyl-terminal fragments of platelet basic protein (PBP) which is found in the alpha-granules of human platelets. NAP-2 is a member of the CXC chemokines. Similar to other ELR domain containing CXC chemokines such as IL-8 and the GRO proteins, NAP-2 has been shown to bind CXCR-2 and to chemoattract and activate neutrophils. Although CTAP-III, β-TG and PBP represent amino-terminal extended variants

of NAP-2 and possess the same CXC chemokine domains, these proteins do not exhibit NAP-2 activity. Recently, it has been shown that the additional amino-terminal residues of CTAP-III masks the critical ELR receptor binding domain that is exposed on NAP-2 and may account for lack of NAP-2 activity.

## **NAP-2/CXCL7 - Protein Information**

## **NAP-2/CXCL7 - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **NAP-2/CXCL7 - Images**