

UBE2C/UBCH10
Catalog # PVGS1221**Specification**

UBE2C/UBCH10 - Product InformationPrimary Accession [O00762](#)**Species**
Human**Sequence**

Met1-Pro179, expressed with additional N-terminal sequence (MHHHHHHAMGIR)

Purity> 95% as analyzed by SDS-PAGE
> 95% as analyzed by HPLC**Endotoxin Level**

< 1 EU/ µg of protein by gel clotting method

Expression System

E. coli

Theoretical Molecular Weight

21.1 kDa

Formulation**Lyophilized from a 0.2 µm filtered solution
in PBS, pH 7.4.****Reconstitution**

It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml.

Storage & Stability

Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

UBE2C/UBCH10 - Additional Information**Gene ID** 11065**Other Names**

Ubiquitin-conjugating enzyme E2 C, 2.3.2.23, (E3-independent) E2 ubiquitin-conjugating enzyme C, 2.3.2.24, E2 ubiquitin-conjugating enzyme C, UbcH10, Ubiquitin carrier protein C, Ubiquitin-protein ligase C, UBE2C, UBCH10

Target Background

Human Ubiquitin Conjugating Enzyme E2 C (UBE2C)/UBCH10 is an essential mediator of mitotic destruction events and cell cycle progression. It catalyzes the destruction of cyclins A and B in

conjunction with the anaphase-promoting complex, and therefore, plays an important role in the control of the cell exit from mitosis. This activity is essential at the end of mitosis for the inactivation of their partner kinase Cdc2 and exit from mitosis into G1 of the next cell cycle. In addition, UBCH10 bears homology to yeast PAS2, a gene that is essential for biogenesis of peroxisomes. UBCH10 is useful for in vitro ubiquitinylation reactions.

UBE2C/UBCH10 - Protein Information

Name UBE2C

Synonyms UBCH10

Function

Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins. In vitro catalyzes 'Lys-11'- and 'Lys-48'-linked polyubiquitination. Acts as an essential factor of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated ubiquitin ligase that controls progression through mitosis. Acts by initiating 'Lys-11'-linked polyubiquitin chains on APC/C substrates, leading to the degradation of APC/C substrates by the proteasome and promoting mitotic exit.

UBE2C/UBCH10 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UBE2C/UBCH10 - Images