

Epigen
Catalog # PVGS1320

Specification

Epigen - Product Information

Primary Accession [O6UW88-2](#)
Species
Human

Sequence
Ala24-Ala95, expressed with an N-terminal Met

Purity
> 95% as analyzed by SDS-PAGE

Endotoxin Level
< 0.2 EU/ µg of protein by gel clotting method

Biological Activity
ED₅₀ < 1.0 µg/ml, measured by a cell proliferation assay using 3T3 cells, corresponding to a specific activity of > 1.0 × 10³ units/mg.

Expression System
E. coli

Formulation **Lyophilized after extensive dialysis against PBS.**

Reconstitution
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH₂O up to 100 µg/ml.

Storage & Stability
Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Epigen - Additional Information

Target Background
Epigen is a cytokine belonging to the Epidermal Growth Factor (EGF) superfamily, which also includes Epiregulin, Amphiregulin, Neuregulin 2-β, and Transforming Growth Factor α. The precursor of Epigen produced in tissues has 154 amino acids, and shares the characteristics of other members of EGF superfamily, including 3 disulfide bonds formed by 6 cysteines. Epigen is present in testis, heart, and liver, and it binds to EGF receptors with a much lower binding affinity than EGF. However, Epigen is more mitogenic than EGF. Epigen achieves its strong mitogenic potency by suppressing ligand-induced receptor inactivation. Unlike EGF, Epigen can also bind to EGF receptors in low pH conditions, helping its recycling. Therefore Epigen has anomalous potency due to its prolonged presence.

Epigen - Protein Information

Epigen - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Epigen - Images