

Betacellulin

Catalog # PVGS1332

Specification

Betacellulin - Product Information

Primary Accession Species Mouse

<u>Q543J8</u>

Sequence Asp32-Tyr111

Purity > 95% as analyzed by SDS-PAGE
> 95% as analyzed by HPLC

Endotoxin Level < 0.2 EU/ μg of protein by gel clotting method

Biological Activity ED₅₀ < 0.08 ng/ml, measured in a cell proliferation assay using 3T3 cells.

Expression System HEK 293

Formulation

Lyophilized after extensive dialysis against PBS.

Reconstitution

It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH₂O or PBS up to 100 μ g/ml.

Storage & Stability

Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Betacellulin - Additional Information

Target Background

Betacellulin (BTC) is a member of the EGF family of growth factors that also includes EGF, TGF- α , Amphiregulin, HB-EGF, Epiregulin, Tomoregulin, Heregulin and Neuregulins. Mature human BTC protein exhibits 80% amino acidsimilarity with mouse BTC protein. BTC is expressed in most tissues including kidney, uterus, liver and pancreas. It is also present in body fluids, including serum, milk, and colostrum. It is synthesized primarily as a transmembrane precursor, which is then processed to a mature molecule by proteolytic events. BTC signals through the EGF receptor.

Betacellulin - Protein Information



Betacellulin - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Betacellulin - Images