

IGF-I
Catalog # PVGS1367

Specification

IGF-I - Product Information

Primary Accession [P05017](#)
Species
Mouse

Sequence
Gly49-Ala118, expressed with an N-terminal Met

Purity
> 95% as analyzed by SDS-PAGE

Endotoxin Level
< 0.2 EU/ µg of protein by gel clotting method

Biological Activity
ED₅₀ < 10.0 ng/ml, measured by a cell proliferation assay using FDC-P1 cells, corresponding to a specific activity of > 1.0 × 10⁵ units/mg.

Expression System
E. coli

Formulation **Lyophilized after extensive dialysis against PBS.**

Reconstitution
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH₂O up to 100 µg/ml.

Storage & Stability
Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

IGF-I - Additional Information

Gene ID 16000

Other Names
Insulin-like growth factor 1 {ECO:0000312|MGI:MGI:96432}, Insulin-like growth factor I, IGF-I, Somatomedin, Igf1 {ECO:0000312|MGI:MGI:96432}

Target Background
Insulin-like Growth Factor I (IGF-I) is a single chain 7 kDa growth-promoting polypeptide originally identified as somatomedin-c. It belongs to the IGF family of peptides, which also includes IGF-II and insulin. The gene expression of IGF-I is mainly regulated by Growth Hormone, and IGF-I executes

its functions via signaling through transmembrane tyrosine receptors (IGF Receptors). Most circulating IGF-I is associated with the IGF Binding Protein 3 (IGFBP-3), and the IGFBPs may inhibit the actions of IGFs by competing against the IGF Receptors. IGF-I is active in post-natal and adult animals, and is crucial for somatic growth, as IGF-I null mice show marked retardation in utero. IGF-I is involved in carcinogenesis, and related to prostate cancer as well.

IGF-I - Protein Information

Name Igf1 {ECO:0000312|MGI:MGI:96432}

Function

The insulin-like growth factors, isolated from plasma, are structurally and functionally related to insulin but have a much higher growth-promoting activity. May be a physiological regulator of [1-14C]- 2-deoxy-D-glucose (2DG) transport and glycogen synthesis in osteoblasts. Stimulates glucose transport in bone-derived osteoblastic (PyMS) cells and is effective at much lower concentrations than insulin, not only regarding glycogen and DNA synthesis but also with regard to enhancing glucose uptake. May play a role in synapse maturation (By similarity).

Ca(2+)-dependent exocytosis of IGF1 is required for sensory perception of smell in the olfactory bulb (PubMed:21496647). Acts as a ligand for IGF1R. Binds to the alpha subunit of IGF1R, leading to the activation of the intrinsic tyrosine kinase activity which autophosphorylates tyrosine residues in the beta subunit thus initiating a cascade of down-stream signaling events leading to activation of the PI3K-AKT/PKB and the Ras-MAPK pathways. Binds to integrins ITGAV:ITGB3 and ITGA6:ITGB4. Its binding to integrins and subsequent ternary complex formation with integrins and IGF1R are essential for IGF1 signaling. Induces the phosphorylation and activation of IGF1R, MAPK3/ERK1, MAPK1/ERK2 and AKT1 (By similarity). As part of the MAPK/ERK signaling pathway, acts as a negative regulator of apoptosis in cardiomyocytes via promotion of STUB1/CHIP-mediated ubiquitination and degradation of ICER-type isoforms of CREM (By similarity).

Cellular Location

Secreted.

IGF-I - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IGF-I - Images