

LIX/CXCL5
Catalog # PVGS1386**Specification**

LIX/CXCL5 - Product Information

Primary Accession [P50228](#)
Species
Mouse

Sequence
Val45-Ala118

Purity
> 98% as analyzed by SDS-PAGE

Endotoxin Level
< 0.2 EU/ µg of protein by gel clotting method

Biological Activity
The EC₅₀ value of mouse LIX/CXCL5 (74aa) on Ca²⁺ mobilization assay in CHO-K1/Gα15/mCXCR2 cells (human Gα15 and mouse CXCR2 stably expressed in CHO-K1 cells) is less than 200.0 ng/ml.

Expression System
HEK 293 Cells

Formulation **Lyophilized after extensive dialysis against PBS.**

Reconstitution
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH₂O or PBS up to 100 µg/ml.

Storage & Stability
Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

LIX/CXCL5 - Additional Information

Gene ID 20311

Other Names
C-X-C motif chemokine 5, Cytokine LIX, Small-inducible cytokine B5, GCP-2(1-78), GCP-2(9-78), Cxcl5, Scyb5

Target Background
Mouse LIX (C-X-C motif chemokine 5) is a small cytokine belonging to the CXC chemokine family that is cleaved into the following 2 chains [GCP-2(1-78) and GCP-2(9-78)]. Mouse LIX plays a role in

reducing sensitivity to sunburn pain in some subjects, and is a potential target which could be used to understand more about pain in other inflammatory conditions. It is most closely related to two highly homologous human neutrophil chemoattractants GCP-2 and ENA-78. The first 78 amino acid residues within the predicted mature mouse LIX shares approximately 61% and 55% amino acid identity with human GCP-2 and ENA-78. This chemokine stimulates the chemotaxis of neutrophils possessing angiogenic properties. It elicits these effects by interacting with the cell surface chemokine receptor CXCR2.

LIX/CXCL5 - Protein Information

Name Cxcl5

Synonyms Scyb5

Function

May participate in the recruitment of inflammatory cells by injured or infected tissue. GCP-2(1-78) and, more potent, GCP-2(9-78) attract neutrophils and are involved in neutrophil activation.

Cellular Location

Secreted.

LIX/CXCL5 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

LIX/CXCL5 - Images