

ENA-78/CXCL5
Catalog # PVGS1389**Specification**

ENA-78/CXCL5 - Product Information

Primary Accession [P42830](#)
Species
Human

Sequence
Ala37-Asn114

Purity
> 98% as analyzed by SDS-PAGE

Endotoxin Level
< 0.2 EU/ µg of protein by gel clotting method

Biological Activity
The EC₅₀ value of human ENA-78/CXCL5 on Ca²⁺ mobilization assay in CHO-K1/ Gα15/hCXCR2 cells (human Gα15 and human CXCR2 stably expressed in CHO-K1 cells) is less than 200.0 ng/ml.

Expression System
HEK 293

Formulation **Lyophilized after extensive dialysis against PBS.**

Reconstitution
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH₂O or PBS up to 100 µg/ml.

Storage & Stability
Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

ENA-78/CXCL5 - Additional Information

Gene ID 6374

Other Names
C-X-C motif chemokine 5, ENA-78(1-78), Epithelial-derived neutrophil-activating protein 78, Neutrophil-activating peptide ENA-78, Small-inducible cytokine B5, ENA-78(8-78), ENA-78(9-78), CXCL5, ENA78, SCYB5

Target Background
Epithelial cell derived neutrophil activating peptide (ENA78) also known as C-X-C motif chemokine 5

(CXCL5), is a small cytokine belonging to the CXC chemokine family. It is produced following stimulation of cells with the inflammatory cytokines interleukin-1 or tumor necrosis factor-alpha. Expression of CXCL5 has also been observed in eosinophils, and can be inhibited with the type II interferon, IFN- γ . This chemokine stimulates the chemotaxis of neutrophils possessing angiogenic properties. Full length CXCL5 (78 aa) is trimmed at the Nterminal end by cathepsin G and chymotrypsin to ENA-74 (74 aa) and ENA-70 (70aa), with the shortened forms showing increased potency relative to full length CXCL5. CXCL5 can signal through the CXCR2 receptor.

ENA-78/CXCL5 - Protein Information

Name CXCL5

Synonyms ENA78, SCYB5

Function

Involved in neutrophil activation. In vitro, ENA-78(8-78) and ENA-78(9-78) show a threefold higher chemotactic activity for neutrophil granulocytes.

Cellular Location

Secreted.

ENA-78/CXCL5 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ENA-78/CXCL5 - Images