

**MIP-2/CXCL2**  
Catalog # PVGS1484**Specification**

---

**MIP-2/CXCL2 - Product Information**Primary Accession [P10889](#)**Species**  
Mouse**Sequence**  
Ala28-Asn100**Purity**  
> 95% as analyzed by SDS-PAGE**Endotoxin Level**  
< 0.2 EU/ µg of protein by gel clotting method**Biological Activity**  
The EC<sub>50</sub> value of Mouse MIP-2/CXCL2 on Ca<sup>2+</sup> mobilization assay in CHO-K1/mCXCR2 (mouse CXCR2 stably expressed in CHO-K1 cells) is less than 200.0 ng/ml.**Expression System**  
HEK 293Formulation **Lyophilized from a 0.2 µm filtered solution in PBS.****Reconstitution**  
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH<sub>2</sub>O or PBS up to 100 µg/ml.**Storage & Stability**  
Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.**MIP-2/CXCL2 - Additional Information****Gene ID** 20310**Other Names**  
C-X-C motif chemokine 2, Macrophage inflammatory protein 2, MIP2, Cxcl2, Mip-2, Mip2, Scyb2**Target Background**  
Chemokine (C-X-C motif) ligand 2 (CXCL2) is a small cytokine belonging to the CXC chemokine family that is also referred to as macrophage inflammatory protein 2-alpha (MIP2-alpha), Growth-regulated protein beta (Gro-beta) and Gro oncogene-2 (Gro-2). CXCL2 is secreted by monocytes and macrophages and is chemotactic for polymorphonuclear leukocytes and

hematopoietic stem cells. CXCL2's amino acid sequence is 90% identical to the amino acid sequence of related chemokine, CXCL1. CXCL2 signals through the CXCR2 receptor.

### **MIP-2/CXCL2 - Protein Information**

**Name** Cxcl2

**Synonyms** Mip-2, Mip2, Scyb2

**Function**

Chemotactic for human polymorphonuclear leukocytes but does not induce chemokinesis or an oxidative burst.

**Cellular Location**

Secreted.

### **MIP-2/CXCL2 - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **MIP-2/CXCL2 - Images**