

MCP-1/CCL2
Catalog # PVGS1498**Specification**

MCP-1/CCL2 - Product Information

Primary Accession [P13500](#)
Species
Human

Sequence
Gln24-Thr99

Purity
> 95% as analyzed by SDS-PAGE

Endotoxin Level
< 0.2 EU/ µg of protein by gel clotting method

Biological Activity
The EC₅₀ value of human MCP-1/CCL2 on Ca²⁺ mobilization assay in CHO-K1/Gα15/hCCR2 cells (human Gα15 and human CCR2 stably expressed in CHO-K1 cells) is less than 1.0 µg/ml.

Expression System
E. coli

Formulation **Lyophilized after extensive dialysis against PBS.**

Reconstitution
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH₂O or PBS up to 100 µg/ml.

Storage & Stability
Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

MCP-1/CCL2 - Additional Information

Gene ID 6347

Other Names
C-C motif chemokine 2, HC11, Monocyte chemoattractant protein 1, Monocyte chemotactic and activating factor, MCAF, Monocyte chemotactic protein 1, MCP-1, Monocyte secretory protein JE, Small-inducible cytokine A2, CCL2, MCP1, SCYA2

Target Background
CCL2, also known as monocyte chemotactic and activating factor (MCAF), was initially purified

independently by two groups based on its ability to chemoattract monocytes. Subsequent to its cloning and sequencing, it became evident that this protein is also identical to the product of the human JE gene. The JE gene, originally identified in mouse fibroblasts, is a platelet-derived growth factor (PDGF) inducible gene. The human CCL2 cDNA encodes a 99 amino acid residue precursor protein with a 23 residue hydrophobic signal peptide that is cleaved to generate the 76 residue mature protein. Natural CCL2 is heterogeneous in size due to the addition of O-linked carbohydrates and sialic acid residues. In addition to fibroblasts, tumor cells, smooth muscle cells, endothelial cells, and mononuclear phagocytes can also produce CCL2 either constitutively or upon stimulation by various stimuli. CCL2 is a member of the β (CC) subfamily of chemokines. Recently, the existence of MCP2 and MCP3 with 62% and 73% amino acid identity respectively, to CCL2 have been reported.

MCP-1/CCL2 - Protein Information

Name CCL2

Synonyms MCP1, SCYA2

Function

Acts as a ligand for C-C chemokine receptor CCR2 (PubMed: [10529171](http://www.uniprot.org/citations/10529171), PubMed: [10587439](http://www.uniprot.org/citations/10587439), PubMed: [9837883](http://www.uniprot.org/citations/9837883)). Signals through binding and activation of CCR2 and induces a strong chemotactic response and mobilization of intracellular calcium ions (PubMed: [10587439](http://www.uniprot.org/citations/10587439), PubMed: [9837883](http://www.uniprot.org/citations/9837883)). Exhibits a chemotactic activity for monocytes and basophils but not neutrophils or eosinophils (PubMed: [8195247](http://www.uniprot.org/citations/8195247), PubMed: [8627182](http://www.uniprot.org/citations/8627182), PubMed: [9792674](http://www.uniprot.org/citations/9792674)). May be involved in the recruitment of monocytes into the arterial wall during the disease process of atherosclerosis (PubMed: [8107690](http://www.uniprot.org/citations/8107690)).

Cellular Location

Secreted

Tissue Location

Expressed in the seminal plasma, endometrial fluid and follicular fluid (at protein level) (PubMed:23765988). Expressed in monocytes (PubMed:2513477).

MCP-1/CCL2 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MCP-1/CCL2 - Images