

GRO- β /CXCL2
Catalog # PVGS1499**Specification**

GRO- β /CXCL2 - Product Information

Primary Accession [P19875](#)
Species
Human

Sequence
Ala35-Asn107

Purity
> 95% as analyzed by SDS-PAGE

Endotoxin Level
< 0.2 EU/ μ g of protein by gel clotting method

Biological Activity
The EC₅₀ value of human GRO β on Ca²⁺ mobilization assay in CHO-K1/G α 15/hCXCR2 cells (human G α 15 and human CXCR2 stably expressed in CHO-K1 cells) is less than 200.0 ng/ml.

Expression System
E. coli

Formulation **Lyophilized after extensive dialysis against PBS.**

Reconstitution
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH₂O or PBS up to 100 μ g/ml.

Storage & Stability
Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

GRO- β /CXCL2 - Additional Information

Gene ID 2920

Other Names
C-X-C motif chemokine 2, Growth-regulated protein beta, Gro-beta, Macrophage inflammatory protein 2-alpha, MIP2-alpha, GRO-beta(5-73), GRO-beta-T, Hematopoietic synergistic factor, HSF, SB-251353, CXCL2, GRO2, GROB, MIP2A, SCYB2

Target Background
Human GRO- α , GRO- β (MIP-2 α), and GRO- γ (MIP-2 β) are products of three distinct,

nonallelic human genes. GRO- β and GRO- γ share 90% and 86% amino acid sequence homology, respectively, with GRO- α . All three isoforms of GRO are CXC chemokines that can signal through the CXCR1 or CXCR2 receptors. GRO expression is inducible by serum or PDGF and/or by a variety of inflammatory mediators, such as IL-1 and TNF, in monocytes, fibroblasts, melanocytes and epithelial cells. In certain tumor cell lines, GRO is expressed constitutively. Similar to other alpha chemokines, the three GRO proteins are potent neutrophil attractants and activators. In addition, these chemokines are also active toward basophils.

GRO- β /CXCL2 - Protein Information

Name CXCL2

Synonyms GRO2, GROB, MIP2A, SCYB2

Function

Produced by activated monocytes and neutrophils and expressed at sites of inflammation. Hematopoietic chemokine, which, in vitro, suppresses hematopoietic progenitor cell proliferation. GRO-beta(5-73) shows a highly enhanced hematopoietic activity.

Cellular Location

Secreted.

GRO- β /CXCL2 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GRO- β /CXCL2 - Images