

**Siglec-2/CD22**  
**Catalog # PVGS1630****Specification**

---

**Siglec-2/CD22 - Product Information**

Primary Accession [P20273](#)  
**Species**  
Human

**Sequence**  
Asp20-Arg687

**Purity**  
> 95% as analyzed by SDS-PAGE<br>> 95% as analyzed by HPLC

**Endotoxin Level**  
≤ 1 EU/ µg of protein by LAL method

**Biological Activity**  
Immobilized Human Siglec2 at 0.5 µg/ml (100 µl/Well). Dose response curve for Biotinylated Anti-Siglec2 Ab with the EC<sub>50</sub> of 0.2 µg/ml determined by ELISA.

**Expression System**  
Expi293

Formulation **Lyophilized from a 0.22 µm filtered solution in PBS, pH 7.4. Normally 5 % trehalose is added as protectant before lyophilization.**

**Reconstitution**  
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in distilled water up to 100 µg/ml.

**Storage & Stability**  
Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Avoid repeated freeze-thaw cycles.

**Siglec-2/CD22 - Additional Information**

**Gene ID** 933

**Other Names**  
B-cell receptor CD22, B-lymphocyte cell adhesion molecule, BL-CAM, Sialic acid-binding Ig-like lectin 2, Siglec-2, T-cell surface antigen Leu-14, CD22, CD22 {ECO:0000303|PubMed:1691828, ECO:0000312|HGNC:HGNC:1643}

**Target Background**  
CD22, or cluster of differentiation-22, is a molecule belonging to the SIGLEC family of lectins. It is found on the surface of mature B cells and to a lesser extent on some immature B cells. CD22 a

member of the immunoglobulin superfamily. CD22 functions as an inhibitory receptor for B cell receptor (BCR) signaling. It is also involved in the B cell trafficking to Peyer's patches in mice.

## **Siglec-2/CD22 - Protein Information**

**Name** CD22 {ECO:0000303|PubMed:1691828, ECO:0000312|HGNC:HGNC:1643}

### **Function**

Most highly expressed siglec (sialic acid-binding immunoglobulin-like lectin) on B-cells that plays a role in various aspects of B-cell biology including differentiation, antigen presentation, and trafficking to bone marrow (PubMed:<a href="http://www.uniprot.org/citations/34330755" target="\_blank">34330755</a>, PubMed:<a href="http://www.uniprot.org/citations/8627166" target="\_blank">8627166</a>). Binds to alpha 2,6-linked sialic acid residues of surface molecules such as CD22 itself, CD45 and IgM in a cis configuration. Can also bind to ligands on other cells as an adhesion molecule in a trans configuration (PubMed:<a href="http://www.uniprot.org/citations/20172905" target="\_blank">20172905</a>). Acts as an inhibitory coreceptor on the surface of B-cells and inhibits B-cell receptor induced signaling, characterized by inhibition of the calcium mobilization and cellular activation. Mechanistically, the immunoreceptor tyrosine-based inhibitory motif domain is phosphorylated by the Src kinase LYN, which in turn leads to the recruitment of the protein tyrosine phosphatase 1/PTPN6, leading to the negative regulation of BCR signaling (PubMed:<a href="http://www.uniprot.org/citations/8627166" target="\_blank">8627166</a>). If this negative signaling from is of sufficient strength, apoptosis of the B-cell can be induced (PubMed:<a href="http://www.uniprot.org/citations/20516366" target="\_blank">20516366</a>).

### **Cellular Location**

Cell membrane; Single-pass type I membrane protein

### **Tissue Location**

B-lymphocytes.

## **Siglec-2/CD22 - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Siglec-2/CD22 - Images**