

**GCP-2/CXCL6**  
**Catalog # PVGS1674****Specification**

---

**GCP-2/CXCL6 - Product Information**

Primary Accession [P80162](#)  
**Species**  
Human

**Sequence**  
Val43-Asn114

**Purity**  
> 98% as analyzed by SDS-PAGE<br>> 98% as analyzed by HPLC

**Endotoxin Level**  
< 1 EU/ µg of protein by LAL method

**Biological Activity**  
Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human neutrophils is in a concentration range of 10.0-50.0 ng/ml.

**Expression System**  
E. coli

**Theoretical Molecular Weight**  
7.9 kDa

Formulation **Lyophilized from a 0.2 µm filtered solution in PBS, pH 7.4.**

**Reconstitution**  
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml.

**Storage & Stability**  
Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

**GCP-2/CXCL6 - Additional Information**

**Gene ID** 6372

**Other Names**  
C-X-C motif chemokine 6, Chemokine alpha 3, CKA-3, Granulocyte chemotactic protein 2, GCP-2, Small-inducible cytokine B6, Small-inducible cytokine B6, N-processed variant 1, Small-inducible cytokine B6, N-processed variant 2, Small-inducible cytokine B6, N-processed variant 3, CXCL6, GCP2, SCYB6

**Target Background**

Granulocyte chemotactic protein 2 (GCP-2) also known as Chemokine (C-X-C motif) ligand 6 (CXCL6) is a small cytokine belonging to the CXC chemokine family. As its former name suggests, GCP-2 is a chemoattractant for neutrophilic granulocytes. Among human CXC chemokines, GCP2 is most closely related to ENA78 (78% amino acid (aa) sequence identity in the mature peptide region and 86% identity in the signal sequence). The structure and sequence of the genes for human GCP2 and ENA78 also exhibit close similarity suggesting the two genes may have originated from a gene duplication. GCP2 can signal through the CXCR1 and CXCR2 receptors.

**GCP-2/CXCL6 - Protein Information**

**Name** CXCL6

**Synonyms** GCP2, SCYB6

**Function**

Chemotactic for neutrophil granulocytes. Signals through binding and activation of its receptors (CXCR1 and CXCR2). In addition to its chemotactic and angiogenic properties, it has strong antibacterial activity against Gram-positive and Gram-negative bacteria (90-fold-higher when compared to CXCL5 and CXCL7).

**Cellular Location**

Secreted.

**GCP-2/CXCL6 - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**GCP-2/CXCL6 - Images**