

**Nectin-4**  
**Catalog # PVGS1705****Specification**

---

**Nectin-4 - Product Information**Primary Accession [XP\\_005541277.1](#)**Species**  
Cynomolgus**Sequence**  
Gly32-Val351**Purity**  
> 95% as determined by Bis-Tris PAGE  
> 95% as determined by HPLC**Endotoxin Level**  
Less than 1EU per µg by the LAL method.**Biological Activity**  
Immobilized Nectin-4, His,Cynomolgus (Cat.No.: Z03827) at 1 µg/ml (100 µl/Well) on the plate can bind Anti-Nectin-4 Antibody.**Expression System**  
HEK293**Theoretical Molecular Weight**  
35.9 kDa**Formulation**  
**Lyophilized from a 0.22 µm filtered solution in PBS, pH 7.4.****Reconstitution**  
Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.**Storage & Stability**  
Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.**Nectin-4 - Additional Information****Target Background**  
Nectin-4 is a tumor-associated antigen. It is probably involved in cell adhesion through trans-homophilic and -heterophilic interactions. Nectin-4 is overexpressed in urothelial carcinoma and several other malignancies. It is a promising target for antibody-drug conjugates (ADCs).**Nectin-4 - Protein Information**

## **Nectin-4 - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Nectin-4 - Images**