

BCMA/TNFRSF17
Catalog # PVGS1729**Specification**

BCMA/TNFRSF17 - Product InformationPrimary Accession [A0A2K5UD97 / F7FFA0](#)**Species**
Cynomolgus/Rhesus macaque**Sequence**
Met1-Ala53**Purity**
> 95% as determined by Bis-Tris PAGE
> 95% as determined by HPLC**Endotoxin Level**
Less than 1EU per µg by the LAL method.**Biological Activity**
Immobilized BCMA/TNFRSF17 hFc Chimera, Cynomolgus/Rhesus macaque (Cat.No.: Z03853) at 1 µg/ml (100 µl/Well) on the plate can bind Biotinylated Anti-BCMA Antibody.**Expression System**
HEK293**Theoretical Molecular Weight**
31.5 kDa**Formulation** **Lyophilized from a 0.22 µm filtered solution in PBS, pH 7.4.****Reconstitution**
Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.**Storage & Stability**
Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.**BCMA/TNFRSF17 - Additional Information****Target Background**
BCMA (B-cell maturation antigen), also called TNFRSF17, is encoded by the TNFRSF17 gene. It acts as a cell surface receptor of the TNF receptor superfamily. BCMA recognizes B-cell activating factor (BAFF). It interacts with the B-cell activating factor TNFSF13B.**BCMA/TNFRSF17 - Protein Information**

BCMA/TNFRSF17 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BCMA/TNFRSF17 - Images