

**BCMA/TNFRSF17**  
**Catalog # PVGS1730****Specification**

---

**BCMA/TNFRSF17 - Product Information**

Primary Accession [O88472-1](#)  
**Species**  
Mouse

**Sequence**  
Met1-Thr49

**Purity**  
> 95% as determined by Bis-Tris PAGE<br/>

**Endotoxin Level**  
Less than 1EU per µg by the LAL method.

**Biological Activity**  
Immobilized BCMA/TNFRSF17, His & Avi, Mouse (Cat.No.: Z03854) at 2 µg/ml (100 µl/Well) on the plate can bind Mouse APRIL, hFc Tag.

**Expression System**  
HEK293

**Theoretical Molecular Weight**  
8.5 kDa

Formulation **Lyophilized from a 0.22 µm filtered solution in PBS, pH 7.4.**

**Reconstitution**  
Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.

**Storage & Stability**  
Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

**BCMA/TNFRSF17 - Additional Information**

**Target Background**  
BCMA (B-cell maturation antigen), also called TNFRSF17, is encoded by the TNFRSF17 gene. It acts as a cell surface receptor of the TNF receptor superfamily. BCMA recognizes B-cell activating factor (BAFF). It interacts with the B-cell activating factor TNFSF13B.

**BCMA/TNFRSF17 - Protein Information**

**BCMA/TNFRSF17 - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**BCMA/TNFRSF17 - Images**