

**MSLN/Mesothelin**  
**Catalog # PVGS1734****Specification**

---

**MSLN/Mesothelin - Product Information**

Primary Accession [Q13421-3](#)  
**Species**  
Human

**Sequence**  
Glu296-Ser598

**Purity**  
> 95% as determined by Bis-Tris PAGE  
> 95% as determined by HPLC

**Endotoxin Level**  
Less than 1EU per µg by the LAL method.

**Biological Activity**  
Immobilized MSLN/Mesothelin(296-598), His, Human (Cat.No.: Z03858) at 1 µg/ml (100 µl/Well) on the plate can bind Biotinylated Human CA125, His Tag.

**Expression System**  
HEK293

**Theoretical Molecular Weight**  
35.23 kDa

Formulation **Lyophilized from a 0.22 µm filtered solution in PBS, pH 7.4.**

**Reconstitution**  
Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.

**Storage & Stability**  
Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

**MSLN/Mesothelin - Additional Information**

**Target Background**  
Mesothelin, also known as MSLN, is encoded by the MSLN gene. It encodes a precursor protein of 71 kDa that is processed to a 31 kDa shed protein called megakaryocyte potentiating factor (MPF) and a 40 kDa fragment, mesothelin. MSLN is attached to the cell membrane by a glycosyl-phosphatidylinositol (GPI) anchor. It is a differentiation antigen which is highly expressed in several human cancers, including virtually all mesotheliomas and pancreatic adenocarcinomas.

## **MSLN/Mesothelin - Protein Information**

## **MSLN/Mesothelin - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **MSLN/Mesothelin - Images**