

FOLR1
Catalog # PVGS1813**Specification**

FOLR1 - Product Information

Primary Accession [P15328](#)
Species
Human

Sequence
Arg25-Met233

Purity
> 95% as determined by Bis-Tris PAGE
> 95% as determined by HPLC

Endotoxin Level
Less than 1EU per µg by the LAL method.

Biological Activity
Immobilized FOLR1, His & Avi, Human (Cat.No.: Z03924) at 1 µg/ml (100 µl/Well) on the plate can bind Anti-FOLR1 Antibody, hFc Tag

Expression System
HEK293

Theoretical Molecular Weight
27.5 kDa

Formulation **Lyophilized from a 0.22 µm filtered solution in PBS, pH 7.4 .**

Reconstitution
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH₂O more than 100 µg/ml.

Storage & Stability
Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

FOLR1 - Additional Information

Gene ID 2348

Other Names
Folate receptor alpha, FR-alpha, Adult folate-binding protein, FBP, Folate receptor 1, Folate receptor, adult, KB cells FBP, Ovarian tumor-associated antigen MOv18, FOLR1, FOLR

Target Background
Folate Receptor 1 (FOLR1), also known as Folate Receptor alpha and Folate Binding Protein (FBP),

is a 37 - 42 kDa protein that mediates the cellular uptake of folic acid and reduced folates. Dietary folates are required for many key metabolic processes including nucleotide and methionine synthesis, the interconversion of glycine and serine, and histidine breakdown. FOLR1 binds to folate and reduced folic acid derivatives and mediates delivery of 5-methyltetrahydrofolate and folate analogs into the interior of cells. Has high affinity for folate and folic acid analogs at neutral pH.

FOLR1 - Protein Information

Name FOLR1

Synonyms FOLR

Function

Binds to folate and reduced folic acid derivatives and mediates delivery of 5-methyltetrahydrofolate and folate analogs into the interior of cells (PubMed:[19074442](http://www.uniprot.org/citations/19074442), PubMed:[23851396](http://www.uniprot.org/citations/23851396), PubMed:[23934049](http://www.uniprot.org/citations/23934049), PubMed:[2527252](http://www.uniprot.org/citations/2527252), PubMed:[8033114](http://www.uniprot.org/citations/8033114), PubMed:[8567728](http://www.uniprot.org/citations/8567728)). Has high affinity for folate and folic acid analogs at neutral pH (PubMed:[23851396](http://www.uniprot.org/citations/23851396), PubMed:[23934049](http://www.uniprot.org/citations/23934049), PubMed:[2527252](http://www.uniprot.org/citations/2527252), PubMed:[8033114](http://www.uniprot.org/citations/8033114), PubMed:[8567728](http://www.uniprot.org/citations/8567728)). Exposure to slightly acidic pH after receptor endocytosis triggers a conformation change that strongly reduces its affinity for folates and mediates their release (PubMed:[8567728](http://www.uniprot.org/citations/8567728)). Required for normal embryonic development and normal cell proliferation (By similarity).

Cellular Location

Cell membrane; Lipid-anchor, GPI-anchor Apical cell membrane; Lipid-anchor, GPI- anchor Basolateral cell membrane; Lipid-anchor, GPI-like-anchor. Secreted Cytoplasmic vesicle. Cytoplasmic vesicle, clathrin-coated vesicle. Endosome. Note=Endocytosed into cytoplasmic vesicles and then recycled to the cell membrane

Tissue Location

Primarily expressed in tissues of epithelial origin. Expression is increased in malignant tissues. Expressed in kidney, lung and cerebellum. Detected in placenta and thymus epithelium.

FOLR1 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

FOLR1 - Images