

# CD3D/CD3 delta

Catalog # PVGS1887

### Specification

# CD3D/CD3 delta - Product Information

Primary Accession Species Human

Sequence Phe22-Ala105

**Purity** > 95% as determined by Bis-Tris PAGE

Endotoxin Level Less than 1EU per  $\mu$ g by the LAL method.

Expression System HEK293

**Theoretical Molecular Weight** 10.4 kDa

Formulation

Reconstitution

Lyophilized from a 0.22 µm filtered solution in PBS[(pH 7.4).

Centrifuge the tube before opening. Reconstituting to a concentration more than 100  $\mu$ g/ml is recommended. Dissolve the lyophilized protein in distilled water.

**Storage & Stability** Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

## CD3D/CD3 delta - Additional Information

Target Background

T-cell surface glycoprotein CD3 delta chain, also known as CD3D, is a single-pass type I membrane protein.In immunology, the CD3 (cluster of differentiation 3) T cell co-receptor helps to activate both the cytotoxic T cell (CD8 naive T cells) and also T helper cells (CD4 naive T cells). It consists of a protein complex and is composed of four distinct chains. In mammals, the complex contains a CD3γ chain, a CD3δ chain, and two CD3ε chains.

### CD3D/CD3 delta - Protein Information

P04234-1



# CD3D/CD3 delta - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

CD3D/CD3 delta - Images