

**Fc gamma RIII/CD16**  
Catalog # PVGS1902**Specification**

---

**Fc gamma RIII/CD16 - Product Information**Primary Accession [Q5D5I8](#)**Species**  
Mouse**Sequence**  
Leu32-Thr215**Purity**  
> 95% as determined by Bis-Tris PAGE  
> 95% as determined by HPLC**Endotoxin Level**  
Less than 1EU per µg by the LAL method.**Biological Activity**  
OKT3 captured on CM5 Chip via Protein A can bind Fc gamma RIII/CD16, His, Mouse in SPR assay (Biacore T200). Test result was comparable to standard batch.**Expression System**  
HEK293**Theoretical Molecular Weight**  
22.2 kDa**Formulation** **Lyophilized from a 0.22 µm filtered solution in PBS, (pH 7.4).****Reconstitution**  
Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.**Storage & Stability**  
Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.**Fc gamma RIII/CD16 - Additional Information****Target Background**  
Immunoglobulin G (IgG) Fc receptors play a critical role in linking IgG antibody-mediated immune responses with cellular effector functions. A high resolution map of the binding site on human IgG1 for human Fc gamma RI, Fc gamma RIIA, Fc gamma RIIB, Fc gamma RIIIA, and FcRn receptors has been determined. A common set of IgG1 residues is involved in binding to all Fc gamma R; Fc gamma RII and Fc gamma RIII also utilize residues outside this common set.

## **Fc gamma RIII/CD16 - Protein Information**

## **Fc gamma RIII/CD16 - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Fc gamma RIII/CD16 - Images**